

## **ScienceDirect**

Journal of the Chinese Medical Association 78 (2015) 377–378



www.jcma-online.com

#### Editorial

# Plant products impairing fertility of animals



Fertility is possibly the most important capability contained within living cells or species. Therefore, finding the most efficacious way to enhance or promote fertility is the main goal of all living things. To best manage nature's strong predisposition for fertility, plant-based traditional medicines for birth control have long been incorporated into medical practices throughout the world, including those in the rural populations of North-East India, China, and Africa. <sup>1–3</sup>

In this issue of the Journal of the Chinese Medical Association, Muhammad and Muhammad<sup>1</sup> published a very interesting review article entitled "Anti-fertility activity of medicinal plants," introducing >60 species of plants that might possess antifertility activity. This article divided these antifertility plants into five categories, namely, plants exhibiting antiovulation activity, anti-implantation activity, antispermatogenic activity, antifertility activity by interfering with the action of hormones, and finally abortifacient activity. The authors claimed to have provided a comprehensive summary of medicinal plants from the literature, including results from Cochrane Central Register of Controlled Trials (CENTRAL), Embase, and PubMed searches using terms such as antianti-implantation, antiovulation, spermatogenic activity of plants. We applaud the authors on the success of this article, because their research and information-gathering task were not easy. However, given the size of the project that they undertook, it goes without saying that their review ultimately will omit certain important data. For example, information about the extract of Drynaria quercifolia (L.) J. Smith rhizome was missed in their review. This extract was proven to possess significant antifertility activity, not only showing a high efficacy for both abortifacient and anti-implantation activities, but also affecting sexhormone release.<sup>2</sup> In addition, the presence of an "adjuvant reagent," including that in solutions or solvents, could influence the final effect of antifertility activity of these medicinal plants. As noted earlier, the methanol extract of D. quercifolia (L.) J. Smith rhizome has shown higher efficacy for both abortifacient (p < 0.01) and anti-implantation performance (p < 0.01) than its aqueous extract dose.<sup>2</sup>

Furthermore, the effect of plant-derived products on antifertility may be dose dependent, reversible, and sex dependent. For example, Shaik and colleagues<sup>4</sup> demonstrated the antifertility potential of methanolic leaf extract of *Artemisia vulgaris*, which showed a strong and significant decrease in implant formation (50% and 100%) at two dose levels, 300 mg/kg and 600 mg/kg, respectively. However, it also manifested a strong estrogenic effect resulting in a significant increase in uterine weight in immature ovariectomized rats. By contrast, although Solomon and coworkers showed that the methanolic root extract of *Rumex steudelii* could cause atrophic changes in the uterus and disruption of ovarian folliculogenesis by inhibiting further development of the recruited ovarian follicles, there was no effect in male albino rats. Dhanapal and coworkers showed the reversible effect of *Enicostemma axillare* leaves and *Urena lobata* root on inhibiting spermatogenesis and steroidogenesis, however, this effect appeared to have no significance in female Wistar albino rats.

Once again, although we commend the publication of this review article, we also conducted an extensive literature review up to December 25, 2014, to investigate the relationship between fertility and plants using the following strategies to identify the publications addressing fertility and plants. The term "decrease, fertility and plants" was used to search PubMed (http://www.ncbi.nlm.nih.gov/pubmed/?term=antifertility%2C+plants), and the term "increase, fertility and plants" in place of the term "decrease, fertility and plants" was used to access relevant articles (http://www.ncbi.nlm.nih.gov/ pubmed/?term=increase%2C+fertility%2C+plants). search criteria identified 266 and 574 published articles, respectively, suggesting that many natural products, especially plants, are still promising targets to be investigated not only for fertility control but also for infertility treatment. This expansive approach is also highly recommended for other troublesome diseases, such as many chronic or aging diseases, including neoplasms, diabetes mellitus, hypertension, and similar maladies. Indeed, paclitaxel, one of the most famous antineoplastic agents extracted from plants, is now widely used in the management of various kinds of neoplastic diseases, including ovarian cancer and breast cancer.

#### **Conflicts of interest**

The authors declare that there are no conflicts of interest related to the subject matter or materials discussed in this article.

### Acknowledgments

This article was supported by grants from the Ministry of Science and Technology, Executive Yuan (Grant No. MOST 103-2314-B-010-043-MY3) and Taipei Veterans General Hospital (Grant Nos. V102C-141; V103C-112; V104C-095; V102E4-003; and V103E4-003).

#### References

- Muhammad D, Muhammad A. Anti-fertility activity of medicinal plants. J Chin Med Assoc 2015;78:382-8.
- Das B, Dey A, Das Talukdar A, Nongalleima Kh, Choudhury MD, Deb L. Antifertility efficacy of *Drynaria quercifolia* (L.) J. Smith on female Wister albino rats. *J Ethnopharmacol* 2014;153:424–9.
- Gebrie E, Makonnen E, Zerihun L, Debella A. The possible mechanisms for the antifertility action of methanolic root extract of *Rumex steudelii*. Afr Health Sci 2005;5:119–25.
- Shaik A, Kanhere RS, Cuddapah R, Nelson KS, Vara PR, Sibyala S. Antifertility activity of *Artemisia vulgaris* leaves on female Wistar rats. Chin J Nat Med 2014;12:180-5.
- Solomon T, Largesse Z, Mekbeb A, Eyasu M, Asfaw D. Effect of Rumex steudelii methanolic root extract on ovarian folliculogenesis and uterine histology in female albino rats. Afr Health Sci 2010;10:353-61.
- Dhanapal R, Ratna JV, Gupta M, Sarathchandran I. Preliminary study on antifertility activity of *Enicostemma axillare* leaves and *Urena lobata* root used in Indian traditional folk medicine. *Asian Pac J Trop Med* 2012;5:616–22.
- Horng HC, Lai CR, Chang WH, Wen KC, Chen YJ, Juang CM, et al. Comparison of early-stage primary serous fallopian tube carcinomas and equivalent stage serous epithelial ovarian carcinomas. *Taiwan J Obstet Gynecol* 2014;53:547–51.
- Huang BS, Sun HD, Hsu YM, Chang WH, Horng HC, Yen MS, et al. Clinical presentation and outcome of adult-type granulosa cell tumors: a retrospective study of 30 patients in a single institute. *J Chin Med Assoc* 2014:77:21-5.

Fa-Kung Lee Department of Obstetrics and Gynecology, Cathay General Hospital, Taipei, Taiwan, ROC Kuan-Hao Tsui Department of Obstetrics and Gynecology, National Yang-Ming University School of Medicine, Taipei, Taiwan, ROC

Department of Obstetrics and Gynecology, Kaohsiung Veterans General Hospital, Kaohsiung, Taiwan, ROC

Department of Pharmacy and Graduate Institute of Pharmaceutical Technology, Tajen University, Pingtung County, Taiwan, ROC

Peng-Hui Wang\* Department of Nursing, National Yang-Ming University School of Nursing, Taipei, Taiwan, ROC

Department of Obstetrics and Gynecology, National Yang-Ming University School of Medicine, Taipei, Taiwan, ROC

Division of Gynecology, Department of Obstetrics and Gynecology, Taipei Veterans General Hospital, Taipei, Taiwan, ROC

Immunology Center, Taipei Veterans General Hospital, Taipei, Taiwan, ROC

Department of Medical Research, China Medical University Hospital, Taichung, Taiwan, ROC

\*Corresponding author. Dr. Peng-Hui Wang, Division of, Gynecology, Department of Obstetrics and Gynecology, Taipei, Veterans General Hospital, 201, Section 2, Shih-Pai Road, Taipei 112, Taiwan, ROC.

E-mail addresses: phwang@vghtpe.gov.tw, phwang@ym.edu.tw (P.-H. Wang)